



# Pre-teens & Teens for Needed Vaccines!

Keep your adolescent patients healthy and safe

Vaccines routinely recommended at 11-12 yrs of age

Tdap

Meningococcal Conjugate

Human Papillomavirus (for girls only)

Catch-up any adolescents that do not have

2 doses of Hepatitis A

2 doses of Var or documentation of immunity\*

Influenza (during flu season)

2 doses of MMR

3 doses of IPV

3 doses of Hep B

Assess for immunizations at EVERY clinic visit-

Look up every child in Michigan Care Improvement Registry (MCIR) before administering vaccines

MCIR provides immunization history, assesses for needed vaccines and checks for minimum age and intervals

Don't miss an opportunity to administer vaccines

Give vaccines at well visits, sick visits, sports and camp physicals

Administer all needed vaccines

\*Criteria for Immunity to Varicella, June 22, 2007

Var should be administered to school age children and adolescents without:

- Documentation of age-appropriate vaccination
  - School-aged children, adolescents, adults: two doses
- Lab evidence of immunity or lab confirmation of disease
- A healthcare provider's: (nurse, NP, physician, PA)
  - Diagnosis of varicella disease or
  - Verification of history of typical varicella disease
- History of herpes zoster (shingles) based on healthcare provider diagnosis

Note: MMRV (ProQuad<sup>®</sup>) is licensed for children 12 months thru 12 years of age only