



ASSESS HIGH SCHOOL STUDENTS FOR NEEDED VACCINES

Assess all your adolescent patients for needed vaccines. Check for:

- ✓ A booster dose of Tdap
- ✓ 1 dose of Meningococcal Conjugate
- ✓ 3 doses of Human Papillomavirus (for girls only)
- ✓ 2 doses of Hepatitis A
- ✓ 2 doses of Var or documentation of immunity*
- ✓ 2 doses of MMR
- ✓ 3 doses of IPV
- ✓ 3 doses of Hep B

Some teens may need additional vaccines either due to their own specific health conditions or exposure in households to other people with age-related or health-related risks. The additional vaccines for which your teenage patients should be assessed for include:

- ✓ Influenza
- ✓ Pneumococcal polysaccharide (PPV23)

*VARICELLA IMMUNITY CRITERIA, JUNE 22, 2007

Varicella vaccine should be administered to school age children and adolescents without:

- Documentation of age-appropriate vaccination
 - School-aged children, adolescents, adults: two doses
- Lab evidence of immunity or lab confirmation of disease
- A healthcare provider's: (nurse, NP, physician, PA)
 - Diagnosis of varicella disease or
 - Verification of history of typical varicella disease
- History of herpes zoster (shingles) based on healthcare provider diagnosis

Note: MMRV (ProQuad[®]) is licensed for children 12 months thru 12 years of age only