

PCV7 Routine and Catch-up Schedule

Age at this visit	Previous PCV7 history	Recommended schedule*
2-6 months	0 doses	3 doses 2 months apart; 4 th dose at 12-15 months
	1 dose	2 doses 2 months apart; 4 th dose at 12-15 months
	2 doses	1 dose, 2 months after the most recent dose; 4 th dose at 12-15 months
7-11 months	0 doses	2 doses 2 months apart; 3 rd dose at 12-15 months
	1 or 2 doses before age 7 months	1 dose at 7-11 months, with another dose at 12-15 months (at least 2 months apart)
12-23 months	0 doses	2 doses at least 2 months apart
	1 dose before age 12 months	2 doses at least 2 months apart
	1 dose at/after 12 months	1 dose at least 2 months after the most recent dose
	2 or 3 doses before age 12 months	1 dose at least 2 months after the most recent dose
24-59 months Healthy children†	Any incomplete schedule	1 dose at least 2 months after the most recent dose
24-59 months High risk§	Any incomplete schedule of less than 3 doses	1 dose least 2 months after the most recent dose and another dose at least 2 months later
	Any incomplete schedule of 3 doses	1 dose at least 2 months after the most recent dose

* For children vaccinated at age less than 1 year, the minimum interval between doses is 4 weeks. Doses given at 12 months of age or older should be spaced at least 8 weeks apart.

† Administer 1 dose of PCV7 to all healthy children aged 24-59 months having any incomplete schedule

§ For children with underlying medical conditions, administer 2 doses of PCV7 at least 8 weeks apart if previously received less than 3 doses or 1 dose of PCV7 if previously received 3 doses. Children with sickle cell disease, asplenia, chronic heart or lung disease, diabetes, cerebrospinal fluid leak, cochlear implant, human immunodeficiency virus infection or another immunocompromising condition, and American Indian or Alaska Native children in areas with a demonstrated risk for invasive pneumococcal disease more than twice the national average (i.e., Alaska, Arizona, New Mexico, and Navajo populations in Colorado and Utah).

References: Preventing Pneumococcal Disease Among Infants and Young Children, MMWR, October 6, 2000, Vol 49, No RR-9